

A NOTE ON CCC FORCINGS

MARCIN SABOK

ABSTRACT. The aim of this short note is to communicate a simple solution to the problem posed in [1] as Question 7.2.7: is it true that for every ccc σ -ideal I any I -positive Borel set contains modulo I an I -positive closed set?

1. INTRODUCTION

We work in a fixed Polish space X with a continuous, strictly positive Borel measure μ . We write $\text{Bor}(X)$ for the family of Borel sets in X .

A σ -ideal in X is a family $I \subseteq \text{Bor}(X)$ which is closed under taking countable unions and subsets. An associated Boolean algebra $\mathbb{P}_I$ is the quotient algebra $\text{Bor}(X)/I$. We say that a σ -ideal I is ccc if any antichain in $\mathbb{P}_I$ is countable.

The most common ccc σ -ideals are: the family of Borel sets of measure zero, denoted by $\mathcal{N}$ and the family of Borel meager sets, denoted by $\mathcal{M}$.

We say that a Borel set B is I -positive if $B \notin I$. It is a well known fact that any $\mathcal{N}$ -positive Borel set contains a $\mathcal{N}$ -positive closed set. On the other hand, for any $\mathcal{M}$ -positive Borel set B there is a closed set C for which $C \setminus B \in \mathcal{M}$, in other words B contains C modulo $\mathcal{M}$. In [1] the author asks if this can be generalized to all ccc σ -ideals.

Question ([1], Question 7.2.7). Suppose that I is a σ -ideal such that $\mathbb{P}_I$ is ccc. Is it true that every positive Borel set contains a positive closed set modulo the ideal I ?

2. THE SOLUTION

We will answer the above question negatively by giving an example of a ccc σ -ideal I and a I -positive Borel set which does not contain modulo I any closed I -positive set.

Take $J = \mathcal{M} \cap \mathcal{N}$. It is clear that J is ccc, so we only need to prove the following proposition.

Proposition. There exists an J -positive Borel set A which does not contain modulo J any J -positive closed set.

Proof. Let A be any Borel set such that $A \in \mathcal{N} \setminus \mathcal{M}$. Such a set can be obtained by decomposing X into two sets $A \in \mathcal{N}$ and $B \in \mathcal{M}$. In particular A is J -positive. Take any closed set C such that $C \setminus A \in J$. Note that this implies that $C \in \mathcal{N}$. But also $C \in \mathcal{M}$ because any set of measure zero must have empty interior. This shows that $C \in J$. $\square$

REFERENCES

- [1] Zapletal J., *Forcing Idealized*, Cambridge Tracts in Mathematics 174, 2008

MATHEMATICAL INSTITUTE, WROCLAW UNIVERSITY, PL. GRUNWALDZKI 2/4,
50-384 WROCLAW, POLAND

E-mail address: `sabok@math.uni.wroc.pl`